


Wisconsin Healthy People 2010 Data Package



A local data package developed by the
Bureau of Health Information

in collaboration with
the Division of Public Health



July, 2003 Updates

Wisconsin Department of Health and Family Services

Crawford County




Leading Health Indicators

Binge Drinking


Adults 18+ who consumed 5 or more drinks on one occasion in the past 30 days

Southern Region



Tobacco Use


Adults 18+ who currently smoke cigarettes
Southern Region



Obesity and Overweight, 2000-2001

Percent of adults 18+ obese and/or overweight based on Body Mass Index (BMI)


Southern Region



Regular and Vigorous Physical Activity


Adults 18+ who exercise 20 minutes or more at 50% of aerobic capacity,
3 or more times per week

Southern Region



Health Insurance Coverage

Adults 18+ with coverage entire preceding year
Southern Region




Mortality

Mortality Due to Coronary Heart Disease

Rates per 100,000 population, age-adjusted to year 2000 standard population


Crawford County



Mortality Due to Cerebrovascular Disease


Rates per 100,000 population, age-adjusted to year 2000 standard population

Crawford County




Mortality Due to Lung Cancer

Rates per 100,000 population, age-adjusted to year 2000 standard population
Crawford County



Mortality Due to Breast Cancer

Rates per 100,000 female population, age-adjusted to year 2000 standard population
Southern Region




Cerebrovascular Disease Deaths, 1997-2001

number of deaths
Crawford County




Lung Cancer Deaths, 1997-2001

Number of deaths
Crawford County




Breast Cancer Deaths, 1997-2001

Number of deaths
Crawford County




Source: Wisconsin
resident death
certificates (BHI,
DHFS)


Coronary Heart Disease Mortality Rates 1997 - 2001



Cerebrovascular Disease Mortality Rates 1997 - 2001



Lung Cancer Mortality Rates 1997 - 2001



Regional Rates

Northeast	47.0
Northern	47.5
Southern	48.3
Southeast	53.9
Western	46.7

State Rate 49.7
2010 Target 44.9

Legend

Rates / 100,000 Population

	31.8 - 45.5
	45.6 - 53.5
	53.6 - 59.4
	59.5 - 86.2

< 50 Deaths (No Rate)


* Rates are age-adjusted to year 2000 standard population

Environmental Factors

Associated with Health


Hospitalizations for Asthma

Rates per 10,000 population, age-adjusted to year 2000 standard population
Crawford County



Children in Medicaid/WIC Screened for Lead


Percent of program-eligible children ever tested
Crawford County



Socioeconomic Indicators

Children In Poverty

Percent of children 0-17 below poverty level, 1999
Crawford County



Poverty: All Ages

Percent of total county population below poverty level, 1999

Crawford County

20%

15%

10%

5%

0%


10.2%

8.7%

Source: U.S.
Census 2000,
long form
sample (U.S.
Census Bureau)

Crawford County


Wisconsin



Median Income

1999 median household income (annual)


Crawford County



Maternal and Child Health

Infant Mortality


Deaths before age one per 1,000 live births
Southern Region



First-Trimester Prenatal Care, 2001


Percent of births to mothers who received early prenatal care

Crawford County



Teen Births, 1997-2001

Number of births to mothers less than 20 years of age
Crawford County




Source: Wisconsin
resident birth certificates
(BHI, DHFS)

Injury and Violence

Motor Vehicle Crashes Related to Alcohol

Percent of crashes where alcohol was a factor
Crawford County




Source: Wisconsin
Department of
Transportation

Homicide rates, 1997-2001 combined

Rates per 100,000 population, age-adjusted to year 2000 standard population

Southern Region



Sexually Transmitted Diseases

Sexually Transmitted Diseases

Number of chlamydia and gonorrhea cases reported, 1997-2001

Crawford County

